

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) EP 1 178 041 A1

(12) EUROPEAN PATENT APPLICATION

(43) Date of publication:  
06.02.2002 Bulletin 2002/06

(51) Int Cl.7: C07D 211/22, A61K 31/445

(21) Application number: 01124314.4

(22) Date of filing: 28.04.1995

(84) Designated Contracting States:  
AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL  
PT SE

(72) Inventor: The designation of the inventor has not yet been filed

(30) Priority: 18.05.1994 US 245731  
11.04.1995 US 417161

(74) Representative: VOSSIUS & PARTNER  
Siebertstrasse 4  
81675 München (DE)


(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:  
95918278.3 / 0 766 668

Remarks:  
This application was filed on 19 - 10 - 2001 as a divisional application to the application mentioned under INID code 62.

(71) Applicant: Aventis Pharmaceuticals Inc.  
Bridgewater, NJ 08807-0800 (US)

(54) Process for preparing anhydrous and hydrate forms of antihistaminic piperidine derivatives


(57) The present invention is related to novel processes for preparing anhydrous and hydrated forms of piperidine derivatives, polymorphs and pseudomorphs thereof of the formulas



which are useful as antihistamines, antiallergic agents and bronchodilators.

EP 1 178 041 A1





wherein

$R_1$  represents hydrogen or hydroxy;

$R_2$  represents hydrogen; or

$R_1$  and  $R_2$  taken together form a second bond between the carbon atoms bearing  $R_1$  and  $R_2$ ;

$n$  is an integer of from 1 to 5;


$R_3$  is  $-\text{CH}_2\text{OH}$ ,  $-\text{COOH}$  or  $-\text{COOalkyl}$  wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;

each of  $A$  is hydrogen or hydroxy; and

pharmaceutically acceptable salts and individual optical isomers thereof,

comprising subjecting the corresponding hydrated, pharmaceutically acceptable acid addition salt to a water-minimizing recrystallization.

**[0004]** In addition, the present invention provides a process for preparing the hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula



wherein

$R_1$  represents hydrogen or hydroxy;

$R_2$  represents hydrogen; or

$R_1$  and  $R_2$  taken together form a second bond between the carbon atoms bearing  $R_1$  and  $R_2$ ;

$n$  is an integer of from 1 to 5;

$R_3$  is  $-\text{CH}_2\text{OH}$ ,  $-\text{COOH}$  or  $-\text{COOalkyl}$  wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;

each of A is hydrogen or hydroxy; and  
pharmaceutically acceptable salts and individual optical isomers thereof,

comprising subjecting the corresponding anhydrous, pharmaceutically acceptable acid addition salts to an aqueous recrystallization.

**[0005]** In addition, the present invention provides processes for preparing polymorphs of anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride designated herein as Form I and Form III and processes for preparing pseudomorphs of hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride designated herein as Form II and Form IV.

**[0006]** The Form I polymorph may be identified by the following characteristics: a visual melting point (capillary tube) in the range of about 196-201°C; a melt endotherm with extrapolated onset in the range of about 195-199°C as determined by differential scanning calorimetry; and an X-ray powder diffraction pattern essentially as shown in **Table 1** wherein the XRPD patterns were measured using a powder diffractometer equipped with a Co X-ray tube source. The sample was illuminated with Co K $\alpha_1$  radiation and XRPD data were collected from 5 to 55° 2 $\theta$ . (intensities may vary radically due to preferred orientation).

Table 1

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
11.8	30
7.3	30
6.3	65
5.9	35
5.0	45
4.8	100
4.4	45
3.9	60
3.8	75
3.7	30

**[0007]** The Form III polymorph may be identified by the following characteristics: a visual melting point (capillary tube) in the range of about 166-171°C; a broad endotherm below about 90°C, a melt endotherm with an extrapolated onset of about 166°C as determined by differential scanning calorimetry; and an X-ray powder diffraction pattern essentially as shown in **Table 2** wherein the XRPD patterns were measured using a powder diffractometer equipped with a Co X-ray tube source. The sample was illuminated with Co K $\alpha_1$  radiation and XRPD data were collected from 5 to 55° 2 $\theta$ . (intensities may vary radically due to preferred orientation).

Table 2

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
9.0	95
4.9	100
4.8	35
4.6	25
4.5	25
3.7	25

**[0008]** The Form II pseudomorph may be identified by the following characteristics: a visual melting point (capillary tube) in the range of about 100-105°C; a large broad endotherm below about 100°C and a small endothermic peak (about 2 joules/gram) with extrapolated onsets in the range of about 124-126°C as determined by differential scanning calorimetry; and an X-ray powder diffraction pattern essentially as shown in **Table 3** wherein the XRPD patterns were measured using a powder diffractometer equipped with a Co X-ray tube source. The sample was illuminated with Co

# EP 1 178 041 A1

K $\alpha_1$  radiation and XRPD data were collected from 5 to 55° 2 $\theta$  (intensities may vary radically due to preferred orientation).

Table 3

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
7.8	45
6.4	44
5.2	85
4.9	60
4.7	80
4.4	55
4.2	50
4.1	60
3.7	75
3.6	60
3.5	50

**[0009]** The Form IV pseudomorph may be identified by the following characteristics: a visual melting point (capillary tube) in the range of about 113-118°C; two broad overlapping endotherms below about 100°C and an additional endotherm with an extrapolated onset at approximately 146°C as determined by differential scanning calorimetry and an X-ray powder diffraction pattern essentially as shown in **Table 4** wherein the XRPD patterns were measured using a powder diffractometer equipped with a Co X-ray tube source. The sample was illuminated with Co K $\alpha_1$  radiation and XRPD data were collected from 5 to 55° 2 $\theta$ . (Intensities may vary radically due to preferred orientation).

Table 4

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
10.4	60
7.0	45
6.4	50
5.3	100
5.2	55
4.3	75
4.1	50
4.0	45
3.8	60
3.5	55

## DETAILED DESCRIPTION OF THE INVENTION

**[0010]** Pharmaceutically acceptable acid addition salts of the compounds of formula (I) and (II), both anhydrous and hydrated, are those of any suitable inorganic or organic acid. Suitable inorganic acids are, for example, hydrochloric, hydrobromic, sulfuric, and phosphoric acids. Suitable organic acids include carboxylic acids, such as, acetic, propionic, glycolic, lactic, pyruvic, malonic, succinic, fumaric, malic, tartaric, citric, cyclamic, ascorbic, maleic, hydroxymaleic, and dihydroxymaleic, benzoic, phenylacetic, 4-aminobenzoic, 4-hydroxybenzoic, anthranilic, cinnamic, salicylic, 4-aminosalicylic, 2-phenoxybenzoic, 2-acetoxybenzoic, and mandelic acid, sulfonic acids, such as, methanesulfonic, ethanesulfonic and  $\beta$ -hydroxyethanesulfonic acid.

**[0011]** As used herein, the term "hydrate" refers to a combination of water with a compound of formula (I) or (II) wherein the water retains its molecular state as water and is either absorbed, adsorbed or contained within a crystal

lattice of the substrate molecule of formula (I) or (II).

**[0012]** As used herein, the term "adsorped" refers to the physical state wherein the water molecule in the hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II) is distributed over the surface of the solid hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II).

**[0013]** As used herein, the term "absorped" refers to the physical state wherein the water molecule in the hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II) is distributed throughout the body of the solid hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II).

**[0014]** Hydrated, pharmaceutically acceptable acid addition salts of the compounds of formula (I) and (II) are those hydrates ranging from essentially 0.10 to 5 molecules of water per molecule of substrate salt of formula (I) or (II).

**[0015]** As used herein, the term "azeotropic mixture" refers to a liquid mixture of two or more substances which behaves like a single substance in that the vapor produced by partial evaporation of liquid has the same composition as the liquid. The constant boiling mixture exhibits either a maximum or minimum boiling point as compared with that of other mixtures of the same substance.

**[0016]** As used herein, the term "azeotropic distillation" refers to a type of distillation in which a substance is added to the mixture to be separated in order to form an azeotropic mixture with one or more of the constituents of the original mixture. The azeotrope or azeotropes thus formed will have boiling points different from the boiling points of the original mixture. As used herein, the term "azeotropic distillation" also refers to co-distillation.

**[0017]** As used herein, the term "water-minimizing recrystallization" refers to a recrystallization wherein the ratio of anhydrous solvent to substrate hydrate is such that the percentage of water present is minimized, thereby inducing precipitation of the anhydrous form of the substrate.

**[0018]** As used herein, the term "aqueous recrystallization" refers to those processes wherein either 1) a solid material is dissolved in a volume of water or a water/organic solvent mixture sufficient to cause dissolution and the solid material recovered by evaporation of the solvent; 2) a solid material is treated with a minimal amount of water or a water/organic solvent mixture which is not sufficient to cause dissolution, heated to obtain dissolution and cooled to induce crystallization or 3) a solid material is dissolved in a volume of water or a water/organic solvent mixture sufficient to cause dissolution and then the solvent is partially evaporated to form a saturated solution which induces crystallization.

**[0019]** As used herein, the term "crystal digestion" refers to that process wherein a solid material is treated with a minimal amount of water or water/organic solvent mixture which is not sufficient to cause dissolution and either heating or stirring at ambient temperature until the desired transformation has taken place.

**[0020]** As used herein, the term "antisolvent" refers to a poor solvent for the substance in question which when added to a solution of the substance, causes the substance to precipitate.

**[0021]** As used herein, the term "suitable temperature" refers to that temperature which is sufficient to cause dissolution and to permit the precipitation of the desired substance either upon addition of an antisolvent or upon removal of the co-solvent by azeotropic distillation.

**[0022]** The anhydrous, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II) may be prepared from the corresponding hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II) by subjecting the corresponding hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II) to an azeotropic distillation.

**[0023]** For example, the appropriate hydrated, pharmaceutically acceptable acid addition salt of piperidine derivatives of the formula (I) and (II) is first dissolved in a volume of a suitable solvent or solvent mixture which is sufficient to cause dissolution. Examples of such solvents are water, C<sub>1</sub>-C<sub>5</sub> alkanols such as methanol, ethanol and the like; ketone solvents such as acetone, methyl ethyl ketone and the like; aliphatic ester solvents such as ethyl acetate, methyl acetate, methyl formate, ethyl formate, isopropyl acetate and the like and aqueous mixtures of these solvents, such as acetone/water, methyl ethyl ketone/water, water/acetone and water/acetone/ethyl acetate. An additional volume of the same solvent used to effect dissolution or second suitable anhydrous antisolvent is then added to this solution, which is then heated to a boiling point which is suitable to azeotropically remove water and other low boiling components. Suitable anhydrous antisolvents for use in the azeotropic distillation are, for example, ketone solvents such as acetone, methyl ethyl ketone and the like; aliphatic ester solvents such as ethyl acetate, methyl acetate, methyl formate, ethyl formate, isopropyl acetate and the like; C<sub>5</sub>-C<sub>8</sub> aliphatic solvents such as pentane, hexane and the like; aliphatic nitriles, such as acetonitrile and mixtures of these solvents such as acetone/ethyl acetate and the like. The azeotropic mixture of water and solvent is removed by distillation until the temperature changes, indicating that the azeotropic mixture is completely removed. The reaction mixture is cooled and the corresponding anhydrous, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II) is recovered from the reaction zone by, for example filtration.

**[0024]** In addition, the anhydrous, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II) may be prepared from the corresponding hydrated, pharmaceutically acceptable acid addition salts

of piperidine derivatives of the formula (I) and (II) by subjecting the corresponding hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II) to a water-minimizing recrystallization.

**[0025]** For example, the appropriate hydrated, pharmaceutically acceptable acid addition salt of piperidine derivatives of the formula (I) and (II) is dissolved in a volume of a suitable anhydrous solvent or solvent mixture which is sufficient to cause dissolution and heated to reflux. Examples of such solvents are water, C<sub>1</sub>-C<sub>5</sub> alkanols such as methanol, ethanol and the like; ketone solvents such as acetone, methyl ethyl ketone and the like; aliphatic ester solvents such as ethyl acetate, methyl acetate, methyl formate, ethyl formate, isopropyl acetate and the like and aqueous mixtures of these solvents, such as acetone/water, methyl ethyl ketone/water, water/acetone and water/acetone/ethyl acetate. An additional volume of the same solvent used to effect dissolution or second suitable anhydrous antisolvent is then added in a quantity sufficient to initiate precipitation of the anhydrous, pharmaceutically acceptable acid addition salt of piperidine derivatives of the formula (I) and (II). Suitable anhydrous antisolvents are, for example, ketone solvents such as acetone, methyl ethyl ketone and the like; aliphatic ester solvents such as ethyl acetate, methyl acetate, methyl formate, ethyl formate, isopropyl acetate and the like; mixtures of ketone solvents and aliphatic ester solvents such as acetone/ethyl acetate and the like; C<sub>5</sub>-C<sub>8</sub> aliphatic solvents such as pentane, hexane and the like; aliphatic nitriles, such as acetonitrile and mixtures of these solvents such as acetone/ethyl acetate and the like as well as mixtures of water and ketone solvents such as acetone/water and the like; and mixtures of water, ketone solvents and aliphatic ester solvents such as acetone/water/ethyl acetate. The reaction mixture is cooled and the corresponding anhydrous, pharmaceutically acceptable acid addition salt of piperidine derivatives of the formula (I) and (II) is recovered from the reaction zone by, for example filtration.

**[0026]** Polymorphic forms of anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Forms I and III) may be prepared by a variety of methods as detailed below.

#### Form III to Form I

**[0027]** For example, anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form I) may be prepared from anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form III), by subjecting the anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form III) to a crystal digestion as described above.

#### Form II to Form III

**[0028]** In addition, anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form III) may be prepared from hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form II), by subjecting the hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form II) to water-minimizing recrystallization as described above.

#### Form II to Form I

**[0029]** In addition, anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form I) may be prepared from hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form II), by subjecting the hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form II) to water-minimizing recrystallization as described above or by subjecting the hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form II) to an azeotropic distillation.

#### Form IV to Form I

**[0030]** In addition, anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form I) may be prepared from hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form IV), by subjecting the hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form IV) to water-minimizing recrystallization or to an azeotropic distillation as described above.

**[0031]** The hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) may be prepared from the corresponding compound of the formula (II) wherein R<sub>3</sub> is -COOalkyl by subjecting the corresponding compound of the formula (II) wherein R<sub>3</sub> is -COOalkyl to a reduction using an appropriate reducing agent, such as sodium borohydride, potassium borohydride, sodium cyanoborohydride, or tetramethylammonium borohydride

In a suitable solvent, such as, methanol, ethanol, isopropyl alcohol or n-butanol, aqueous mixtures thereof or basic solutions thereof, at temperatures ranging from about 0°C to the reflux temperature of the solvent, and the reaction time varies from about 1/2 hour to 8 hours. After quenching and acidifying with an suitable acid, such as hydrochloric acid, the hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) are recovered from the reaction zone by crystallization and filtration.

**[0032]** In addition, the hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II) may be prepared from the corresponding anhydrous, pharmaceutically acceptable acid addition salts of the formula (I) and (II) by subjecting the corresponding anhydrous, pharmaceutically acceptable acid addition salts of formula (I) and (II) to an aqueous recrystallization.

**[0033]** For example, the appropriate anhydrous, pharmaceutically acceptable acid addition salt of piperidine derivatives of the formula (I) and (II) is treated with a minimal volume of water or suitable water/organic solvent mixture which is insufficient to cause dissolution and heated to reflux. The reaction mixture is cooled and the corresponding hydrated, pharmaceutically acceptable acid addition salt of piperidine derivatives of the formula (I) and (II) is recovered from the reaction zone by, for example filtration. Alternatively, the appropriate anhydrous, pharmaceutically acceptable acid addition salt of piperidine derivatives of the formula (I) and (II) is treated with a volume of water or a suitable water/organic solvent mixture which is sufficient to cause dissolution and the water or water/organic solvent is partially or completely evaporated to a volume which induces crystallization of the hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II). Suitable solvents for use in the above recrystallization are water, acetone/water, ethanol/water, methyl ethyl ketone/aqueous methanol, methyl ethyl ketone/water and the like.

**[0034]** The pseudomorphic forms of hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Forms II and IV) may be prepared by a variety of methods as detailed below.

#### Ethyl Ester/Ketone to Form II

**[0035]** Hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form IV) may be prepared from ethyl 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- $\alpha,\alpha$ -dimethylbenzeneacetate, hydrochloride or free base as described above for the general preparation of the hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) from the corresponding compound of the formula (II) wherein  $R_3$  is -COOalkyl, but rapidly adding water over a period of time ranging from 1 minute to 45 minutes at a temperature range of about -20°C to 50°C to precipitate the hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form II).

#### Ethyl Ester/Ketone to Form IV

**[0036]** Hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form IV) may be prepared from ethyl 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- $\alpha,\alpha$ -dimethylbenzeneacetate, hydrochloride or free base as described above for the general preparation of the hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) from the corresponding compound of the formula (II) wherein  $R_3$  is -COOalkyl, but slowly adding water over a period of time ranging from about 30 minutes to 24 hours and at a temperature range of about 0°C to 50°C, optionally with seeding, to precipitate the hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form IV).

#### Form I to Form II

**[0037]** Hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form II) may be prepared from anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form I) by subjecting hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form II) to an aqueous recrystallization as defined above.

**[0038]** Starting materials for use in the present invention are readily available to one of ordinary skill in the art. For example, ethyl 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- $\alpha,\alpha$ -dimethylbenzeneacetate, hydrochloride is described in U.S. Patent 4,254,129, March 3, 1981.

**[0039]** The following examples present typical processes for preparing the anhydrous and hydrated, pharmaceutically acceptable acid addition salts of piperidine derivatives of the formula (I) and (II), polymorphs and pseudomorphs thereof. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the following terms have the indicated meanings: "g" refers to grams; "mol" refers to mole; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "mp" refers to melting point;



"°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

[0040] Differential Scanning Calorimetry analysis were performed using a TA 2910 DSC with open aluminum pans. The samples were heated to 240°C at 5°C/minute with a 50mL/minute nitrogen purge.

[0041] X-Ray Powder Diffraction analyses were performed as follows:

[0042] The samples were loaded into a quartz (zero scatter) sample holder for the XRPD pattern measurement. The XRPD patterns were measured using a powder diffractometer equipped with a Co X-ray tube source, primary beam monochromator, and position sensitive detector (PSD). The incident beam was collimated using a 1° divergence slit. The active area on the PSD subtended approximately 5°2θ. The source was operated at 35 kV and 30 mA and the sample was illuminated with Co Kα<sub>1</sub> radiation. XRPD data were collected from 5 to 55° 2θ at a rate of 0.25°2θ/minute and a step width of 0.02°2θ. The XRPD patterns were measured without the addition of an internal calibrant.

[0043] Peak positions and Intensities for the most prominent features were measured using a double-derivative peak picking method. X-ray peaks with I/I<sub>0</sub> greater than 20% were reported. The cutoff was chosen arbitrarily. The intensities are rounded to the nearest 5%. Certain peaks appear sensitive to preferred orientation that is caused by changes in crystallite morphology. This results in large changes in the I/I<sub>0</sub> value.

#### Example 1-Preparation of Form II

##### 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]-α,α-dimethylbenzeneacetic acid hydrochloride

#### Method A

[0044] Mix ethyl 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]-α,α-dimethylbenzeneacetate, hydrochloride (101.92g, 0.1807mol) and methanol (510mL) and stir. Rapidly add 50% sodium hydroxide (72.27g, 0.903mol) and wash in with water (61mL). Heat to reflux for 2 hours, allow to cool to 35°C and treat with sodium borohydride (3.42g, 0.0903mol). Add water (100mL) and maintain at 35°C for 10 hours. Add 37% hydrochloric acid (53.0g) to adjust pH to 11.5. Add acetone (26.5mL) and water (102mL). Hold at 35°C for 2 hours and adjust to pH 2.5 with 37% hydrochloric acid (44.69g). Dilute with water (408mL), cool to -15°C, stir for 1.5 hours and collect the precipitate by vacuum filtration. Wash the filtercake with deionized water (3X100mL) and vacuum dry to give 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]-α,α-dimethylbenzeneacetic acid hydrochloride hydrate (97.10g).

#### Method B

[0045] Place ethyl 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]-α,α-dimethylbenzeneacetate, hydrochloride (60.01g, 0.106mol) in a 1-L three necked round-bottom flask and fit the flask with a mechanical stirrer, a Claisen head, a thermometer and a reflux condenser with a nitrogen bubbler on top. Add methanol (300mL) and turn the stirrer on. Dilute the slurry with water (60mL) and heat to 52-54°C over 15-20 minutes. Hold at 52°C for 2 hours and then add 50% sodium hydroxide (42.54g, 0.532mol). Heat at 73°C for approximately 1 hour, 45 minutes, cool to less than 35°C using a water bath and then add sodium borohydride (2.02g, 0.0534mol). Stir overnight at 35°C, treat with acetone (15.5 mL) and stir for 2 hours at 35°C. Acidify the mixture to a pH of 1.85 with 28% hydrochloric acid (75.72g), dilute with water (282 mL), stir for about 30 minutes and cool over about 2 hours to -15°C. Filter the solids off and wash with water (2X75mL) and ethyl acetate (2X75mL). Vacuum dry the solid and allow to stand for 2 days to give 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]-α,α-dimethylbenzeneacetic acid hydrochloride hydrate (Form II) (57.97g, 91.5%) as a fine powder.

#### Method C

[0046] Place ethyl 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]-α,α-dimethylbenzeneacetate (56.12g, 0.1064mol) in a 1-L three necked round-bottom flask and fit the flask with a mechanical stirrer, a Claisen head, a thermometer and a reflux condenser with a nitrogen bubbler on top. Add methanol (300mL) and turn the stirrer on. Dilute the slurry with water (60mL) and heat to reflux using a heating mantle controlled by a Therm-O-Watch. When the mixture reaches about 35°C, treat with 50% sodium hydroxide (34.05g, 0.4256mol) and rinse in with water (42mL). Stir at reflux for 2 hours, 15 minutes, cool over 1 hour to 35°C and then treat with sodium borohydride (2.02g, 0.0534mol). Stir for 7.5 hours and allow to stand at room temperature without stirring for 1.75 days. Warm the mixture to 35°C and quench with acetone (15.5mL, 0.21mol) and stir for 2 hours. Add water (60mL) and adjust the pH to 2.5 with 32% hydrochloric acid (65.22g). Cool to 40°C and rinse the pH probe with water (25mL). Add water over about 30 minutes (192mL), hold the temperature at 33°C for 10 minutes and add a few seed crystals. Cool the slurry to -12°C over about 45 minutes and isolate the solid by filtration (586.2g). Wash with water (2X100mL) and then with ethyl

acetate (100mL, prechilled to about -10°C). Vacuum dry overnight (1 mmHg, 50°C) to give 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form II) (58.86g, 98%) as a white solid.

#### 5 Example 2-Preparation of Form IV

4-[4-[4-(hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form IV)

10 **[0047]** Place ethyl 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidiny]-1-oxobutyl]- $\alpha,\alpha$ -dimethylbenzeneacetate (56.12g, 0.1064mol) in a 1-L three necked round-bottom flask and fit the flask with a mechanical stirrer, a Claisen head, a thermometer and a reflux condenser with a nitrogen bubbler on top. Add methanol (300mL) and turn the stirrer on. Dilute the slurry with water (60mL) and heat to reflux using a heating mantle controlled by a Therm-O-Watch. When the mixture reaches about 35°C, treat with 50% sodium hydroxide (34.05g, 0.4256mol) and rinse in with water (42mL).  
 15 Stir at reflux for 2 hours, 15 minutes, cool over 1 hour to 35°C and then treat with sodium borohydride (2.02g, 0.0534mol). Stir for 7.5 hours and allow to stand at room temperature without stirring for 1.75 days. Warm the mixture to 35°C and quench with acetone (15.5mL, 0.21mol) and stir for 2 hours. Add water (60mL) and adjust the pH to 2.5 with 32% hydrochloric acid (65.22g). Cool to 40°C and rinse the pH probe with water (25mL). Hold the temperature at 33°C for 10 minutes, add a few seed crystals and add water over about 4 hours (192mL) at 35°C. Cool the slurry to  
 20 -12°C over about 45 minutes and isolate the solid by filtration (586.2g). Wash with water (2X100mL) and then with ethyl acetate (100mL, prechilled to about -10°C). Vacuum dry overnight (1 mmHg, 50°C) to give 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form IV); mp 115-116°C (dec).

25 XRPD: **Table 5**

**[0048]**

Table 5

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
10.38	60
6.97	45
6.41	50
5.55	30
5.32	100
5.23	55
5.11	35
4.98	25
4.64	30
4.32	35
4.28	75
4.12	50
4.02	45
3.83	60
3.65	20
3.51	55
3.46	25
2.83	20

Example 3-Conversion of Form II to Form I4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form I)

5 **[0049]** Treat 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form II) (20.0g, 0.0355mol) with deionized water (2g) and add acetone (60mL) in small portions over several minutes with stirring. Filter through filter aid and wash the filter cake with acetone (30mL). Wash the filtercake with acetone (22mL), reflux filtrate and then slowly add ethyl acetate (32mL over 15 minutes) keeping the mixture at reflux. Reflux for 10 minutes, then slowly add additional ethyl acetate (23mL over 10 minutes) and reflux for an additional  
10 15 minutes. Add additional ethyl acetate (60mL over 5-10 minutes) and continue refluxing for 15 minutes. Cool to approximately 8°C in an ice bath, filter the solid and wash with ethyl acetate (85mL). Vacuum dry at 55°C for 1.5 hours to give the title compound (18.16g, 95%).

Example 4-Conversion of Form II to Form I4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochlorideMethod A:

20 **[0050]** Treat 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form II) (5.00g, 0.0083mol) with methylethyl ketone (130mL). Slowly add water (0.4mL), filter through filter aid and wash the filter cake with methylethyl ketone (20mL). Heat to reflux and distill off 75mL of solvent, cool to -15°C and collect by vacuum filtration. Wash with methylethyl ketone (2X10mL) and vacuum dry at 60°C to give the title compound (4.33g, 97%); mp 196-198°C.

Method B:

30 **[0051]** Treat 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form II) (1.4g) with acetone (60mL) and heat to reflux. Reduce the volume to approximately 35mL to remove all water which boils off as an azeotrope (88/12:acetone/water). Cool the solution and collect the title compound as a crystalline solid.

Method C:

35 **[0052]** Mix 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form II) (53.88g, 0.100mol) and add water (4.79g) and methyl ethyl ketone (240mL). Stir until the solid is slurried up and add additional methyl ethyl ketone (1L). Stir for 0.5 hours, filter through a pad of filter aid, wash the filtercake with methyl ethyl ketone (100mL) and transfer the filtrate and wash to a 2L, 3-necked flask fitted with a thermometer, mechanical stirrer and distillation head. Distill off a total of 721mL of methyl ethyl ketone, cool and stir  
40 over 1 hour to 40°C. Cool to -15°C and hold for 10 minutes. Collect the solid by vacuum filtration and wash the filtercake with methyl ethyl ketone (2X65mL) and vacuum dry at 55°C overnight to give the title compound (52.76g, 97.9%); mp 197.5-200°C.

Method D:

45 **[0053]** Treat 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form II) (40.0g, 0.0696mol, assayed at 93.6% purity, having 0.89g water present and 35.1g, 0.0575mol, assayed at 88.0% purity, having 2.47g water present) with water (8.30g; the amount calculated to bring the weight of water present to 17% of the anhydrous weight of 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate, taking into account the water in the hydrated salt). Add methyl ethyl ketone (approximately 500mL) and stir until most of the solids dissolve. Add additional methyl ethyl ketone (700mL) in portions over approximately 10 minutes and continue stirring for 1/2 hour. Filter through a thin pad of filter aid, wash the filtercake and flask with additional methyl ethyl ketone (100mL) and transfer to a boiling flask fitted with a thermometer, mechanical stirrer, heating mantle, a 12-plate Oldershaw (vacuum-jacketed) distillation column and a distillation head with the capability of regulating the reflux ratio in a rough fashion, washing in with additional  
55 methyl ethyl ketone (100mL). Distill off 450 mL of solvent, cool to -15°C and filter the solid. Wash with methyl ethyl ketone (2X100mL) and dry to give the title compound (68.3g, 99.9%); mp 197-199°C.

Method E

**[0054]** Bring methyl ethyl ketone (4mL) to a boil and add 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (500mg). Decant the top layer and add methyl ethyl ketone (3mL) to the aqueous layer. Boil the solution until the temperature reached 79°C, reduce the volume by 25%, remove from heat and cover with aluminum foil. Allow the solution to cool, filter the resulting crystals and air dry to give the title compound.

Example 5-Conversion of Form I to Form II

4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate

Method A

**[0055]** Treat 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form I) (2.0g) with ethanol (4mL) and deionized water (20mL). Heat at 80°C until a solution is formed and then stir at room temperature for 23 hours. Filter the resulting slurry, wash with water (2X10mL) and dry under vacuum at 35°C overnight to give the title compound (1.88g); mp 100-105°C.

XRPD: **Table 6**

**[0056]**

Table 6

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
11.41	20
7.98	20
7.83	45
6.58	45
6.42	60
5.66	20
5.52	45
5.39	30
5.23	65
5.14	45
4.86	65
4.72	100
4.45	65
4.40	45
4.32	45
4.18	45
4.06	65
4.02	55
3.85	25
3.79	75
3.74	95

Table 6 (continued)

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
3.61	80
3.56	25
3.47	65
3.41	20
2.74	20

**Method B**

**[0057]** Mix water (35.5mL), methanol (26.3mL) and sodium chloride (2.59g). Add 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form I) (4.77g). Heat to reflux on a steam bath until dissolution and cool to -10°C. Filter the resulting solid, wash with water (2X25mL) and vacuum dry overnight to give the title compound (4.80g).

**Example 6-Conversion of Form II into Form III**

4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form III)

**[0058]** Place 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form II) (55.56g, 0.0929mol having 10% water) in a pressure bottle along with water (2.96g) and acetone (38.1g). Seal the bottle tightly and heat to approximately 80°C. Cool to about 50°C, filter through filter aid in a coarse sintered glass funnel and dilute with acetone (90g). Transfer to a 1L flask fitted with a mechanical stirrer, thermometer and a reflux condenser. Heat the mixture to reflux and allow to cool and stir over the weekend. Cool to -15°C and filter on a coarse sintered glass funnel, wash with ethyl acetate (2X50mL) and vacuum dry at 50°C.

**[0059]** Place a majority of the solid obtained (45.24g) in a 500 mL three necked flask fitted with a mechanical stirrer, thermometer and a reflux condenser. Add acetone (240mL) and water (4.82g) and reflux the mixture overnight. Allow the slurry to cool to 35°C and place in an ice water bath and cool to less than 5°C. Filter the solid off on a coarse sintered glass funnel, wash with ethyl acetate (50mL) and vacuum dry at 50°C for several hours to give the title compound as a white crystalline powder (43.83g, 97%); mp 166.5-170.5°C.

XRPD: Table 7

**[0060]**

Table 7

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
8.95	95
4.99	20
4.88	100
4.75	35
4.57	25
4.47	25
4.46	20
3.67	20
3.65	25

Example 7-Conversion of Form III Into Form I

4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form I)]

**[0061]** Place 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form III) (40.0g as an ethyl acetate wetcake-27.9g dry basis) in a 1L three necked flask fitted with a mechanical stirrer, thermometer and a reflux condenser. Add acetone (240mL) and heat the mixture to reflux for about 20 hours. Cool the slurry to -15°C and isolate the solids by filtration on a coarse sintered glass frit funnel. Wash with ethyl acetate (50mL) and vacuum dry overnight to give the title compound (26.1g, 93.7%); mp 197.5-199.5°C.

XRPD: **Table 8**

**[0062]**

Table 8

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
11.75	35
7.23	35
6.24	60
5.89	40
5.02	20
4.94	30
4.83	100
4.44	30
3.93	75
3.83	20
3.77	85
3.71	25
3.62	30
3.32	25
3.31	20

Example 8-Conversion of Form IV into Form I

4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride (Form I)]

**[0063]** Place 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form IV) (54.35g, 0.0970mol, having 4% water present) in a pressure bottle along with water (4.16g) and acetone (38.1g). Seal the bottle tightly and heat to approximately 80°C. Cool to less than 60°C, filter through filter aid in a coarse sintered glass funnel and rinse the filter cake with acetone (32.4g). Place acetone (215g) in a 1L three necked flask fitted with a mechanical stirrer, thermometer, a reflux condenser and containing a small amount of Form I crystals and heat to reflux. Add a portion of the acetone/water solution of 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form IV) (47.65g) to the refluxing acetone over about 10 minutes. Slowly add ethyl acetate (157.5g) over 45 minutes then add the remaining portion of the acetone/water solution of 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride hydrate (Form IV), rinsed in with about 20mL of acetone. Add additional ethyl acetate (157.5g) over 45 minutes to 1 hour, maintaining the slurry at reflux. Stir for 15 minutes, cool to -15°C and vacuum filter the white solid on a 350mL coarse sintered glass funnel. Wash the solids with ethyl acetate (2X50mL) and vacuum dry overnight to give the title compound (50.36g, 97%); mp 198-199.5°C.

XRPD: Table 9

[0064]

Table 9

D-Space, Angstroms	Intensity, I/I <sub>0</sub> , %
14.89	20
11.85	20
7.30	20
6.28	70
5.91	25
5.55	20
5.05	25
4.96	55
4.85	100
4.57	45
4.45	55
3.94	45
3.89	20
3.84	20
3.78	60
3.72	35
3.63	20
3.07	20
3.04	20
2.45	20

[0065] The polymorphic and pseudomorphic 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride compounds of this invention are useful as antihistamines, antiallergy agents and bronchodilators and may be administered alone or with suitable pharmaceutical carriers, and can be in solid or liquid form such as, tablets, capsules, powders, solutions, suspensions or emulsions.

[0066] The polymorphic and pseudomorphic 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride compounds of this invention can be administered orally, parenterally, for example, subcutaneously, intravenously, intramuscularly, intraperitoneally, by intranasal instillation or by application to mucous membranes, such as, that of the nose, throat and bronchial tubes, for example, in an aerosol spray containing small particles of a compound of this invention in a spray or dry powder form.

[0067] The quantity of polymorphic or pseudomorphic 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride compound administered will vary depending on the patient and the mode of administration and can be any effective amount. The quantity of polymorphic or pseudomorphic 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride compound administered may vary over a wide range to provide in a unit dosage an effective amount of from about 0.01 to 20 mg/kg of body weight of the patient per day to achieve the desired effect. For example, the desired antihistamine, antiallergy and bronchodilator effects can be obtained by consumption of a unit dosage form such as a tablet containing 1 to 500 mg of a polymorphic or pseudomorphic 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride compound of this invention taken 1 to 4 times daily.

[0068] The solid unit dosage forms can be of the conventional type. Thus, the solid form can be a capsule which can be the ordinary gelatin type containing a polymorphic or pseudomorphic 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride compound of this invention and a carrier, for example,

lubricants and inert fillers such as lactose, sucrose or comstarch. In another embodiment the polymorphic or pseudomorph 4-[4-(4-(hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride compound is tableted with conventional tablet bases such as lactose, sucrose or comstarch or gelatin, disintegrating agents such as comstarch, potato starch or alginic acid, and a lubricant such as stearic acid or magnesium stearate.


**[0069]** The polymorphic or pseudomorph 4-[4-(4-(hydroxydiphenylmethyl)-1-piperidiny]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride compounds of this invention may also be administered in injectable dosages by solution or suspension of the compounds in a physiologically acceptable diluent with a pharmaceutical carrier which can be a sterile liquid such as water and oils, with or without the addition of a surfactant and other pharmaceutically acceptable adjuvants. Illustrative of oils there can be mentioned those of petroleum, animal, vegetable or synthetic origin, for example, peanut oil, soybean oil or mineral oil. In general, water, saline, aqueous dextrose and related sugar solutions and glycols such as propylene glycol or polyethylene glycol are preferred liquid carriers, particularly for injectable solutions.

**[0070]** For use as aerosols the compounds of this invention in solution or suspension may be packaged in a pressurized aerosol container together with suitable propellants, for example, hydrocarbon propellants such as, propane, butane or isobutane with the usual adjuvants as may be administered in a non-pressurized form such as in a nebulizer or atomizer.

**[0071]** The term patient as used herein is taken to mean warm blooded animals, birds, mammals, for example, humans, cats, dogs, horses, sheep, bovine cows, pigs, lambs, rats, mice and guinea pigs.

## Claims

1. A process for preparing a compound of the formula



wherein

R<sub>1</sub> represents hydrogen or hydroxy;

R<sub>2</sub> represents hydrogen; or

R<sub>1</sub> and R<sub>2</sub> taken together form a second bond between the carbon atoms bearing R<sub>1</sub> and R<sub>2</sub>;

n is an integer of from 1 to 5;

R<sub>3</sub> is -CH<sub>2</sub>OH, -COOH or -COOalkyl wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;

W is -CH(OH)- or -C(=O)-;


A is hydrogen or hydroxy;

Y is a pharmaceutically acceptable acid; and

the individual optical isomers thereof,


comprising subjecting a compound of the formula





and the individual optical isomers thereof, wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, n, W, A and Y are defined above and X is a number ranging essentially from 0.10 to 5 to an azeotropic distillation.

2. A process for preparing a compound of the formula



wherein

R<sub>1</sub> represents hydrogen or hydroxy;

R<sub>2</sub> represents hydrogen; or

R<sub>1</sub> and R<sub>2</sub> taken together form a second bond between the carbon atoms bearing R<sub>1</sub> and R<sub>2</sub>;

n is an integer of from 1 to 5;

R<sub>3</sub> is -CH<sub>2</sub>OH, -COOH or -COOalkyl wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;

W is -CH(OH)- or -C(=O)-;


A is hydrogen or hydroxy;

Y is a pharmaceutically acceptable acid;

X is a number ranging essentially from 0.1 to 5; and


the individual optical isomers thereof,

comprising subjecting a compound of the formula




and the individual optical isomers thereof wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $n$ ,  $A$ ,  $W$  and  $Y$  are as defined above, to an aqueous recrystallization.

3. A process for preparing a compound of the formula




and the individual optical isomers thereof, wherein  $Y$  is a pharmaceutically acceptable acid, comprising subjecting a compound of the formula




wherein Y is as defined above and X is a number ranging essentially from 0.10 to 5 to an azeotropic distillation.

4. A process for preparing a compound of the formula




and the individual optical isomers thereof, wherein Y is a pharmaceutically acceptable acid and X is a number ranging essentially from 0.10 to 5, comprising subjecting a compound of the formula




wherein Y is as defined above to an aqueous recrystallization.

5. A process according to Claim 3 or 4 wherein Y is HCl.
6. A process according to Claim 5 wherein x is a number ranging from 1 to 4.
7. A process according to Claim 6 wherein x is a number ranging from 2 to 3.
8. A compound of the formula



wherein X is a number ranging from 0.1 to 5 and the individual optical isomers thereof, preparable by the process comprising the steps of:

a) treating a compound of the formula



and the individual optical isomers thereof with a suitable solvent;

b) heating the mixture to a suitable temperature; and

c) cooling the mixture to initiate crystallization.

9. Form I anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]-α,α-dimethylbenzeneacetic acid hydrochloride,

Form II hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]-α,α-dimethylbenzeneacetic acid hydrochloride,

Form III anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]-α,α-dimethylbenzeneacetic acid hydrochloride and

Form IV hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]-α,α-dimethylbenzeneacetic acid hydrochloride.

10. A pharmaceutical composition comprising an a compound of Claim 9 in admixture or otherwise in association with an inert carrier.

11. Use of a compound as defined in claim 9 for the preparation of a pharmaceutical composition for treating allergic

reactions.

12. A process for preparing the Form I anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ ,  
 $\alpha$ -dimethylbenzeneacetic acid hydrochloride which comprises subjecting Form II hydrated 4-[4-[4-(Hydroxydiphenyl-  
methyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ , $\alpha$ -dimethylbenzeneacetic acid hydrochloride to an azeotropic distillation.
13. A process for preparing the Form I anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ ,  
 $\alpha$ -dimethylbenzeneacetic acid hydrochloride which comprises subjecting Form III anhydrous 4-[4-[4-(Hydroxydiphe-  
nylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ , $\alpha$ -dimethylbenzeneacetic acid hydrochloride to a crystal digestion.
14. A process for preparing the Form I anhydrous 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ ,  
 $\alpha$ -dimethylbenzeneacetic acid hydrochloride which comprises subjecting Form IV hydrated 4-[4-[4-(Hydroxydiphe-  
nylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ , $\alpha$ -dimethylbenzeneacetic acid hydrochloride to an azeotropic distilla-  
tion.
15. A process according to claim 14 wherein suitable solvents or solvent mixtures which are sufficient to cause dis-  
solution are methanol, acetone/water and methyl ethyl ketone/water and a suitable anhydrous antisolvent is methyl  
ethyl ketone.
16. A process for preparing the Form II hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ ,  
 $\alpha$ -dimethylbenzeneacetic acid hydrochloride which comprises:
  - a) reacting ethyl 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- $\alpha$ , $\alpha$ -dimethylbenzeneacetate hy-  
drochloride with an appropriate reducing agent in a suitable solvent;
  - b) acidifying with hydrochloric acid; and
  - c) adding water over a period of time ranging from 1 minute to 45 minutes at a temperature range of about  
-20°C to 50°C.
17. A process for preparing the Form II hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ ,  
 $\alpha$ -dimethylbenzeneacetic acid hydrochloride which comprises:
  - a) reacting ethyl 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- $\alpha$ , $\alpha$ -dimethylbenzeneacetate with  
an appropriate reducing agent in a suitable solvent;
  - b) acidifying with hydrochloric acid; and
  - c) adding water over a period of time ranging from 1 minute to 45 minutes at a temperature range of about  
-20°C to 50°C.
18. A process according to claim 17 wherein the water is added over a period of time ranging from 10 minutes to 30  
minutes at a temperature range of about 0°C to 40°C.
19. A process according to claim 18 wherein the water is added over a period of time ranging from 10 minutes to 30  
minutes at a temperature range of about 20°C to 40°C.
20. A process for preparing the Form II hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ , $\alpha$ -  
 $\alpha$ -dimethylbenzeneacetic acid hydrochloride which comprises subjecting Form I anhydrous 4-[4-[4-(Hydroxydiphenyl-  
methyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ , $\alpha$ -dimethylbenzeneacetic acid hydrochloride to an aqueous recrystalliza-  
tion.
21. A process for preparing the Form IV hydrated 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha$ , $\alpha$ -  
 $\alpha$ -dimethylbenzeneacetic acid hydrochloride which comprises:
  - a) reacting ethyl 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- $\alpha$ , $\alpha$ -dimethylbenzeneacetate hy-  
drochloride with an appropriate reducing agent in a suitable solvent;

b) acidifying with hydrochloric acid; and

22. A process for preparing the Form IV hydrated 4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- $\alpha,\alpha$ -dimethylbenzeneacetic acid hydrochloride which comprises:

5

a) reacting ethyl 4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- $\alpha,\alpha$ -dimethylbenzeneacetate with an appropriate reducing agent in a suitable solvent;

10

b) acidifying with hydrochloric acid; and

c) adding water over a period of time ranging from 30 minutes to 24 hours at a temperature range of about 0°C to 50°C.

15

23. A process according to claim 22 wherein the water is added over a period of time ranging from 1 hour to 12 hours at a temperature range of about 10°C to 40°C.

24. A process according to claim 23 wherein the water is added over a period of time ranging from 2 hours to 8 hours at a temperature range of about 20°C to 40°C.

20

25

30

35

40

45

50

55



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 01 12 4314

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	WO 93 21156 A (MERRELL DOW PHARM. INC.) 28 October 1993 (1993-10-28) Title compound of step e on page 34 * page 1, line 12 *	8-24	C07D211/22 A61K31/445
E	WO 96 26726 A (HOECHST MARION ROUSSEL INC) 6 September 1996 (1996-09-06) * claim 5 *	8	
X	US 4 285 957 A (CARR) 25 August 1981 (1981-08-25) Examples 1-3	8-24	
D, X	US 4 254 129 A (CARR) 3 March 1981 (1981-03-03) Examples 1-3	8-24	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C07D A61K
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 12 December 2001	Examiner Gettins, M
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date O : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technical background O : non-written disclosure P : intermediate document			

EPO FORM 1505 (3.62 P-94) (2001)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 12 4314

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

12-12-2001

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9321156	A	28-10-1993	AU 3797393 A	18-11-1993
			EP 0635004 A1	25-01-1995
			FI 944713 A	02-12-1994
			FI 20002460 A	09-11-2000
			HU 70517 A2	30-10-1995
			JP 8502022 T	05-03-1996
			KR 257805 B1	01-07-2000
			MX 9302076 A1	31-03-1994
			NO 943794 A	09-12-1994
			NO 981431 A	09-12-1994
			NO 981432 A	09-12-1994
			NO 981433 A	09-12-1994
			NO 981434 A	09-12-1994
			NZ 251132 A	20-12-1996
			WO 9321156 A1	28-10-1993
			US 5631375 A	20-05-1997
			US 5675009 A	07-10-1997
			US 5644061 A	01-07-1997
			US 5618940 A	08-04-1997
			US 5652370 A	29-07-1997
			US 5654433 A	05-08-1997
			US 5650516 A	22-07-1997
			US 5663353 A	02-09-1997
			ZA 9302454 A	20-10-1993
WO 9626726	A	06-09-1996	AU 701042 B2	21-01-1999
			AU 4909896 A	18-09-1996
			CA 2213700 A1	06-09-1996
			CN 1176599 A	18-03-1998
			EP 0812195 A1	17-12-1997
			FI 973518 A	27-08-1997
			HU 9802086 A2	28-01-1999
			JP 11501028 T	26-01-1999
			NO 973938 A	28-10-1997
			NZ 302926 A	28-10-1998
			WO 9626726 A1	06-09-1996
			US 6113942 A	05-09-2000
			US 5738872 A	14-04-1998
			US 2001022973 A1	20-09-2001
			US 5855912 A	05-01-1999
US 4285957	A	25-08-1981	US 4254129 A	03-03-1981
			AT 376208 B	25-10-1984
			AT 144880 A	15-03-1984

LHC FORIA PAGES

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82



**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 12 4314

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for those particulars which are merely given for the purpose of information.

12-12-2001

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4285957	A		AU 531146 B2	11-08-1983
			AU 5501680 A	16-10-1980
			BE 882703 A1	31-07-1980
			CA 1123438 A1	11-05-1982
			CH 643245 A5	30-05-1984
			DE 3007498 A1	23-10-1980
			DK 132980 A ,B,	11-10-1980
			ES 489934 D0	16-03-1981
			ES 8103735 A1	16-06-1981
			FR 2453854 A1	07-11-1980
			GB 2048258 A ,B	10-12-1980
			IE 49672 B1	27-11-1985
			IL 59158 A	30-04-1984
			IT 1143960 B	29-10-1986
			JP 1032823 B	10-07-1989
			JP 1555761 C	23-04-1990
			JP 55141469 A	05-11-1980
			NL 8000754 A ,B,	14-10-1980
			NO 801014 A ,B,	13-10-1980
			NZ 192653 A	16-03-1982
			PH 17023 A	11-05-1984
			SE 448726 B	16-03-1987
			SE 8002634 A	11-10-1980
			ZA 8000332 A	28-01-1981
US 4254129	A	03-03-1981	AT 376208 B	25-10-1984
			AT 144880 A	15-03-1984
			AU 531146 B2	11-08-1983
			AU 5501680 A	16-10-1980
			BE 882703 A1	31-07-1980
			CA 1123438 A1	11-05-1982
			CH 643245 A5	30-05-1984
			DE 3007498 A1	23-10-1980
			DK 132980 A ,B,	11-10-1980
			ES 489934 D0	16-03-1981
			ES 8103735 A1	16-06-1981
			FR 2453854 A1	07-11-1980
			GB 2048258 A ,B	10-12-1980
			IE 49672 B1	27-11-1985
			IL 59158 A	30-04-1984
			IT 1143960 B	29-10-1986
			JP 1032823 B	10-07-1989
			JP 1555761 C	23-04-1990
			JP 55141469 A	05-11-1980
			NL 8000754 A ,B,	14-10-1980
			NO 801014 A ,B,	13-10-1980

EPO FORM P0001

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 12 4314

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

12-12-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4254129 A	NZ	192653 A	16-03-1982
	PH	17023 A	11-05-1984
	SE	448726 B	16-03-1987
	SE	8002634 A	11-10-1980
	US	4285957 A	25-08-1981
	ZA	8000332 A	28-01-1981
<hr/>			

EPO FORM 1045/9

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**